

High level of heavy metals in crab meat

Crab meat is a popular seafood, but it sometimes contains large amounts of environmental toxins. The content is so high in many places in Norway that consumption of brown crab meat should generally be discouraged.

Maria Averina

Maria Averina, specialist in clinical chemistry, senior consultant at the Department of Laboratory Medicine, University Hospital of North Norway, and associate professor at UiT The Arctic University of Norway. She is a board member in the special interest group for environmental toxins and public health. The author has completed the ICMJE form and declares no conflicts of interest.

Anne-Lise Bjørke-Monsen almo@helse-bergen.no

Anne-Lise Bjørke-Monsen, specialist in paediatrics and clinical chemistry. She is a senior consultant at the Blood Centre and Clinical Chemistry Department, Innlandet Hospital, and at the Department for Clinical Chemistry and Pharmacology, Haukeland University Hospital, and a board member in the special interest group for environmental toxins and public health. The author has completed the ICMJE form and declares no conflicts of interest.

Bjørn J. Bolann

Bjørn J. Bolann, specialist in clinical chemistry, professor emeritus at the Department of Clinical Science, University of Bergen, and retired senior consultant from the Department of Clinical Chemistry and Pharmacology, Haukeland University Hospital. He is a board member in the special interest group for environmental toxins and public health. The author has completed the ICMJE form and declares no conflicts of interest.

Jan Brox

Jan Brox, specialist in clinical chemistry and head senior consultant at the Department of Laboratory Medicine, University Hospital of North Norway. He is a board member in the special interest group for environmental toxins and public health. The author has completed the ICMJE form and declares no conflicts of interest.

Merete Eggesbø

Merete Eggesbø, PhD, and senior researcher at the Division of Climate and Environmental Health, Norwegian Institute of Public Health, and associate professor at NTNU.

She is a board member in the special interest group for environmental toxins and public health. The author has completed the ICMJE form and declares no conflicts of interest.

Ingrid Hokstad


Ingrid Hokstad, specialty registrar in clinical chemistry at Innlandet Hospital. She is a board member in the special interest group for environmental toxins and public health. The author has completed the ICMJE form and declares no conflicts of interest.

Sandra Huber

Sandra Huber, special advisor at the Department of Laboratory Medicine, University Hospital of North Norway. She is a board member in the special interest group for environmental toxins and public health. The author has completed the ICMJE form and declares no conflicts of interest.

Peter Ørebech

Peter Ørebech, professor in law at the Norwegian College of Fishery Science, UiT The Arctic University of Norway. He is a board member in the special interest group for environmental toxins and public health. The author has completed the ICMJE form and declares no conflicts of interest.



Illustrations: Børge Bredebekk/byHandstekst

Clinical laboratories sometimes receive enquiries about heavy metal poisoning following the consumption of crab meat. Patients' symptoms are often diffuse, and can include pain in the extremities, paraesthesia, fatigue and mood swings. Many patients also have elevated levels of heavy metals, such as mercury, cadmium and lead. It is not clear whether the heavy metals explain the symptoms, but they are potent neurotoxins capable of causing the symptoms described (1).

Patients' symptoms are often diffuse, and can include pain in the extremities, paraesthesia, fatigue and mood swings

It has become increasingly clear that all levels of heavy metals are harmful to living organisms. They can have carcinogenic effects, and cause damage to most organs, even in minute quantities (1). The limits for acceptable intake of mercury, cadmium and lead have therefore been lowered significantly over the past 50 years (2–4). For many metals, there are still not established intake thresholds or clinical diagnostic cut-off points. One such example is tin, which is found in seafood (5), and is neurotoxic and harmful to the liver (6).

In line with international agreements, emissions of heavy metals have been reduced (7). Between 1995 and 2019, emissions of cadmium and mercury were reduced by 80%, and arsenic by 35% (7). Emissions of lead to air have also been significantly reduced, though ammunition remains a considerable source of lead in soil (7).

Nevertheless, there are still large accumulations of both inorganic and organic pollutants in Norwegian fjords, where the concentrations of heavy metals are 2–4 times higher than in Norway's coastal waters (8). The Sørfjorden in Hardanger is an example of the extensive pollution caused by industrial activity. Here the levels of cadmium, lead, mercury, dioxins and polychlorinated biphenyls are so high that the recommendation is not to eat crab, shellfish or fish from parts of the fjord. The Norwegian Food Safety Authority (NFSA) keeps a list of areas where the consumption of certain types of seafood is not recommended (9).

Environmental toxins in crab meat

Consumption of white crab meat, crab roe and brown crab meat is common in Norway. Brown crab meat mainly consists of the crab's hepatopancreas, where most of the environmental toxins are found. The European Food Safety Authority (EFSA) has set a maximum concentration for cadmium of 0.5 mg/kg wet weight for white crab meat, but no such limit has yet been set for brown crab meat (10). The Norwegian Institute of Marine Research found that some crab paste and stuffed crab shells purchased from Norwegian grocery stores in 2016 contained up to 1.8 mg and 1.4 mg of cadmium per kilo respectively (11).

There has been a significant increase in toxic metals in brown crab meat in recent years. Between 2016 and 2018, the average level of cadmium increased by 137%

Ervik et al. examined crabs caught in the Mausundvær archipelago in Trøndelag, and show that there has been a significant increase in toxic metals in brown crab meat in recent years (12, 13). Between 2016 and 2018, the average level of cadmium increased by 137%, lead by 22%, mercury by 133%, copper by 22%, and arsenic by 31%. In 2018, brown crab meat had a median cadmium concentration of 11.9 mg/kg dry weight (3.7 mg/kg wet weight) and a maximum concentration of 202 mg/kg dry weight (63.1 mg/kg wet weight), which is 7 and 126 times higher respectively than the EFSA's maximum allowed concentration of 0.5 mg/kg wet weight in white crab meat (13).

Where does the pollution come from??

The increase in heavy metals over a relatively short period of time indicates that crabs are exposed to ongoing and increasing pollution. It has been difficult to explain this, as emissions of various toxic elements have generally been reduced in Norway. The crabs in the study by Ervik et al. were caught at various locations in one of Norway's main aquaculture areas (12, 13). There are ten fish farms in this area, the oldest of which has been in operation since 1996.

In 2018, a total of 1.35 million tonnes of farmed salmon and trout (wet weight) were sold in Norway, which required 1.8 million tonnes of fish feed (dry weight) (14). The fish feed contains both inorganic and organic pollutants. The environmental toxins that are not deposited in the farmed fish are left at the bottom of the fjord under open farming cages, in the form of uneaten feed and fish faeces.

Table 1 shows measurements of arsenic, lead, cadmium and mercury in fish feed, farmed salmon and trout, taken by the NFSA and the Norwegian Institute of Marine Research (Seafood Data) in 2018 (15, 16). Emissions from fish farming are calculated as content in fish feed minus content in produced salmon and trout. After comparing this with data from the Norwegian Environment Agency on total emissions in Norway (7), we calculated the share that comes from aquaculture. The aquaculture industry is not listed as a source of emissions of these substances on the Norwegian Environment Agency's website. The environment status page on the website specifically states that emissions of cadmium from fish feed are not included in the calculation of total emissions (7). If we include the emissions from aquaculture, total emissions of cadmium are an incredible 29 per cent higher (Table 1).

Table 1 Emissions of arsenic and some heavy metals in Norway in 2018

	Arsenic (As)	Cadmium (Cd)	Mercury (Hg)	Lead (Pb)
<i>Emissions from aquaculture</i>				
Mean concentration in fish feed in 2018 (mg/kg) (15)	2.4	0.16	0.03	0.04

Content in 1.8 million tonnes (consumption 2018) (kg) (15)	4 320	288	54	72
Mean concentration in farmed salmon and trout, 2018 (16)	0.89 mg/kg	< LOQ ₁	0.02 mg/kg	< LOQ ₁
Content in 1.35 million tonnes of salmon and trout (production figures 2018) (kg)	1 202	-	27	-
Emissions: content in fish feed minus content in produced salmon and trout (kg)	3 118	288	27	72
<i>Total emissions in Norway</i>				
Emissions in Norway in accordance with the Norwegian Environment Agency in 2019 (7)	23 tonnes	1 000 kg	500 kg	87 tonnes
<i>Addition from aquaculture in 2018 (%)</i>	+ 14	+ 29	+ 6	+ < 0.1

¹ LOQ: Limit of quantification

The emissions from uneaten feed and fish faeces are spread or accumulated depending on local currents and will impact on the facilities to varying degrees. In 2013, the Norwegian Environment Agency examined cadmium and copper levels in crabs caught at a distance of 80–1000 metres from three salmon farms in Sogn og Fjordane, Møre og Romsdal and Nord-Trøndelag respectively (17). Aquaculture uses around 1700 tonnes of copper annually to impregnate farming nets (figures from 2019), and around 80–90% of this leaks into the sea and accumulates on the seabed (18).

The study concluded that there were no clear indications that fish farms are a source of cadmium in the marine environment. However, at two of the farms, the mean level of cadmium and copper was 1.6–2.4 times higher in brown crab meat caught 80 metres from the farm compared to 1000 metres. At the third facility, the level of cadmium and copper was the same regardless of the distance to the facility. At all locations, the levels of cadmium and copper were co-variants (Spearman correlation coefficient 0.9–1.0), which suggests that they stem from a common source.

Toxic elements will accumulate in fjords where farming takes place

Arsenic, lead, cadmium and mercury are natural elements and as such do not expire. As salmon and trout are farmed in an ongoing process with only a three-month break between each production cycle, toxic elements will accumulate in fjords where farming takes place.

Are crabs edible?

In several places in Norway, crabs are so contaminated that the NFSA advises against all consumption (19). It is clear that many Norwegian fjords have a significant pollution problem. The studies by Ervik et al. show that in just six years there has been a considerable increase in dangerous environmental toxins in crabs caught at Mausundvær (12, 13), which is particularly worrying.

The NFSA currently has several recommendations for how much crab meat different groups in the population can eat based on threshold values for various environmental toxins, but these only apply to white crab meat. The absence of a threshold value for brown crab meat does not, however, mean that its environmental toxins are not dangerous. The NFSA therefore recommends that pregnant women do not eat brown crab meat (20). However, the content of environmental toxins in brown crab meat is so high that we believe this recommendation should apply to everyone.

The accumulation of environmental toxins in crabs and other marine life is a marker of the state of our fjords' health. The increasing toxic content in crab meat is a sign that developments are going in the wrong direction. Aquaculture's contribution to this is significant (Table 1).

If this development is allowed to continue, all seafood from the fjords will become a risk to health and inedible

If this development is allowed to continue, all seafood from the fjords will become a risk to health and inedible. Neither healthcare personnel nor the general public seem to be aware of this problem. We are calling for measures to control these emissions. We must demand that those doing the polluting clean up after themselves, and that emissions are discontinued. Protecting life in the fjords means protecting life for us all.

This article was written on behalf of the special interest group for environmental toxins and public health.

Received 29 April 2022, first revision submitted 13 May 2022, accepted 9 August 2022.

References

1. Balali-Mood M, Naseri K, Tahergorabi Z et al. Toxic Mechanisms of Five Heavy Metals: Mercury, Lead, Chromium, Cadmium, and Arsenic. *Front Pharmacol* 2021; 12: 643972.
2. Ye BJ, Kim BG, Jeon MJ et al. Evaluation of mercury exposure level, clinical diagnosis and treatment for mercury intoxication. *Ann Occup Environ Med* 2016; 28: 5.
3. Satarug S, Haswell-Elkins MR, Moore MR. Safe levels of cadmium intake to prevent renal toxicity in human subjects. *Br J Nutr* 2000; 84: 791–802.
4. Wani AL, Ara A, Usmani JA. Lead toxicity: a review. *Interdiscip Toxicol* 2015; 8: 55–64.
5. Hannisdal R, Bernhard A, Lunestad BT et al. Monitoring program for pharmaceuticals, illegal substances, and contaminants in farmed fish – Annual report for 2018. Havforskningsinstituttet, 2019. https://www.mattilsynet.no/mat_og_vann/produksjon_av_mat/fisk_og_sjomat/oppdrettsfisk/aarsrapport_2018_medisinrester_i_oppdrettsfisk.36741/binary/%C3%85rsrapport%202018%20Medisinrester%20i%20oppdrettsfisk Lest 9.8.2022.
6. Winship KA. Toxicity of tin and its compounds. *Adverse Drug React Acute Poisoning Rev* 1988; 7: 19–38.
7. Miljødirektoratet. Miljøgifter og andre prioriterte stoffer. <https://miljostatus.miljodirektoratet.no/tema/miljogifter/prioritertemiljogifter/> Lest 17.2.2022.
8. Everaert G, Ruus A, Hjermand DØ et al. Additive Models Reveal Sources of Metals and Organic Pollutants in Norwegian Marine Sediments. *Environ Sci Technol* 2017; 51: 12764–73.
9. Mattilsynet. Oversikt over havner, fjorder og innsjøer med forurensning. https://www.matportalen.no/matvaregrupper/tema/fisk_og_skalldyr/oversikt_over_havner_fjorder_og_innsjoer_med_forurensning Lest 17.2.2022.
10. European Food Safety Authority Cadmium in food - Scientific opinion of the Panel on Contaminants in the Food Chain. *EFSA J* 2009; 7: 980.
11. Frantzen S, Duinker A, Sanden M et al. Kadmianalyser i produkter av taskekrabbe kjøpt inn i 2016. Havforskningsinstituttet, 2018. https://www.hi.no/hi/nettrapporter/rapport-fra-havforskningen/2018/kadmianalyser_i_produkter_av_taskekrabbe Lest 8.3.2022.
12. Ervik H, Finne TE, Jenssen BM. Toxic and essential elements in seafood from Mausund, Norway. *Environ Sci Pollut Res Int* 2018; 25: 7409–17.
13. Ervik H, Lierhagen S, Asimakopoulou AG. Elemental content of brown crab (*Cancer pagurus*) - Is it safe for human consumption? A recent case study from Mausund, Norway. *Sci Total Environ* 2020; 716: 135175.
14. Fauske M. Nøkkeltall fra norsk havbruksnæring 2018. Rapport. Fiskeridirektoratet, 2019. <https://fdir.brage.unit.no/fdir-xmlui/bitstream/handle/11250/2826560/nokkeltall-havbruk-2018.pdf?sequence=1&isAllowed=y> Lest 9.8.2022
15. Sele V, Sanden M, Berntssen MHG et al. Program for overvåking av fiskefor. Årsrapport for prøver innsamlet i 2018. Havforskningen, 2019. https://www.mattilsynet.no/dyr_og_dyrhold/for/rapport_overvaakingsprogram_fiskefor_2018.35512/binary/Rapport:%20Overv%C3%A5kingsprogram%20fiskefor%C3%B4r%202018 Lest 9.8.2022.
16. Havforskningsinstituttet. Sjømatdata. <https://sjomatdata.hi.no> Lest 9.8.2022.
17. Falk AH. Kadmium rundt oppdrettsanlegg – Er fiskefor en potensiell kilde til kadmium i marint miljø? Rapport nr. 6676-01. Akvaplan-niva, 2014. <https://www.statsforvalteren.no/siteassets/fm-nordland/dokument-fmno/miljo-og-klima-dokumenter/vann/kadmium-rundt-oppdrettsanlegg-2014---akvaplan-niva-rapport-6676-01.pdf> Lest 9.8.2022
18. Havforskningsinstituttet. Høyt kobberforbruk i oppdrettsnæringen. <https://www.hi.no/hi/nyheter/2021/februar/hoytkobberforbruk-i-oppdrettsnæringen> Lest 17.2.2022.
19. Mattilsynet. Ikke spis selvfanger krabbe fra Salten. https://www.matportalen.no/matvaregrupper/tema/fisk_og_skalldyr/ikke_spis_selvfanger_krabbe_fra_salten Lest 17.2.2022.

20. Mattilsynet. Barn, gravide og ammende bør ikke spise brun krabbemat. https://www.matportalen.no/matvaregrupper/tema/fisk_og_skalldyr/barn_gravide_og_ammende_bor_ikke_spise_brun_krabbemat Lest 18.2.2022.